



**UNIVERSITI PUTRA MALAYSIA**

**BASAL STEM ROT OF OIL PALM :  
INCIDENCE, ETIOLOGY AND CONTROL**

**KHAIRUDIN BIN HASHIM**

**FP 1990 5**

**BASAL STEM ROT OF OIL PALM :  
INCIDENCE, ETIOLOGY AND CONTROL**

By

**KHAIRUDIN BIN HASHIM**

Thesis Submitted in Fulfilment of the  
Requirements for the Degree of Master of Agricultural  
Science in the Faculty of Agriculture  
Universiti Pertanian Malaysia

July 1990



## **ACKNOWLEDGEMENTS**

The author is deeply indebted to his supervisor, Associate Professor Dr. Lim Tong Kwee and co-supervisor, Associate Professor Dr. Abdul Rahman Abdul Razak, for their guidance, invaluable advice and constant encouragement throughout the duration of this study and in the preparation of this thesis.

He is similarly grateful to Mr. Teoh Cheng Hai, Director of Research, Research and Development Department and Messrs Golden Hope Plantations Berhad, for the encouragement, permission and financial support in making the pursuit of this graduate programme possible.

Appreciation is also extended to colleagues and staff of Prang Besar Research Station and Oil Palm Research Station. In particular, I would like to thank Messrs. Poon Yew Chin, Mohamad Hashim bin Ahmad Tajudin and Ho Cheng Tuck for their assistance and useful comments; Messrs. Mohd. Yazair bin Razali and Azman bin Shafie for assistance in the conduct of trials; Miss M. Mageswary and Miss S. Sivaranchitham for typing the early and final drafts of the thesis.

Thanks are also due to the Managers of the estates for the excellent cooperation extended during the assessment on Basal Stem Rot, and to the Agrochemical companies for supplying the test fungicides.

## TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS .....	ii
LIST OF TABLES .....	viii
LIST OF FIGURES .....	xi
LIST OF PLATES .....	xii
ABSTRACT .....	xvi
ABSTRAK .....	xx
CHAPTER	
I INTRODUCTION .....	1
II LITERATURE REVIEW .....	7
Occurrence of Basal Stem Rot .....	7
Symptoms of Basal Stem Rot .....	9
Taxonomy and Nomenclature of Basal Stem Rot Causal Agent .....	10
Growth of <u>Ganoderma</u> spp. <u>In Vitro</u> .....	12
Induction of Sporophore .....	13
Pathogenicity of Basal Stem Rot Pathogen .....	14
Control of Basal Stem Rot .....	16
Cultural Practices .....	16



	<b>Page</b>
Resistance Varieties .....	17
Chemical Control .....	17
III ASSESSMENT ON BASAL STEM ROT INCIDENCE .....	19
Materials and Methods .....	19
Results .....	23
Incidence of Basal Stem Rot .....	23
Quantification of Losses .....	25
Influence of Previous Crops on Basal Stem Rot .....	31
Influence of Replanting Techniques .....	36
Influence of Soil Types .....	38
Influence of Age of Planting Materials .....	39
Discussion .....	41
IV CULTURE AND IDENTIFICATION OF <u>GANODERMA</u> SPP. .	46
Materials and Methods .....	46
Isolation and Preparation of <u>Ganoderma</u> Spawn .....	46
Effect of Substrates for Mycelial Colonisation and Sporophore Induction .....	50
Comparison of <u>Ganoderma</u> Isolates on Standard Substrate .....	57
Results .....	59

	Page
Effect of Substrates for Mycelial Colonisation and Sporophore Induction .....	59
Mycelial Colonisation .....	59
Sporophore Induction .....	62
Comparison of <u>Ganoderma</u> Isolates on Standard Substrate .....	66
Comparative Growth Rate of Isolates .....	66
Comparison of Sporophores .....	70
Comparison of Basidiospores .....	79
Discussion .....	84
V    PATHOGENICITY OF <u>GANODERMA BONINENSE</u> .....	93
Materials and Methods .....	93
Results .....	102
Discussion .....	112
VI   CHEMICAL CONTROL .....	117
Materials and Methods .....	117
<u>In Vitro</u> Screening of Systemic Fungicides .....	117
<u>In Vivo</u> Fungicide Screening .....	119
Results .....	124
<u>In Vitro</u> Screening of Systemic Fungicides .....	124

	<b>Page</b>
<u>In Vivo</u> Fungicide Screening .....	126
Discussion .....	131
VII SUMMARY AND CONCLUSION .....	134
BIBLIOGRAPHY .....	142
APPENDICES .....	147
A Fertilizer Application Schedule and Composition .....	148
B Analysis of Variance for Bunch Counts and Foliar Status at 27 Months from Trial Commencement .....	150
VITA .....	151





## LIST OF TABLES

Table	Page
1 Incidence of Basal Stem Rot in 16 Estates .....	26
2 Census on Palms Lost and Cost for Control in 16 Estates for 1987 .....	28
3 Census on Palms Lost and Cost for Control in 16 Estates for 1988 .....	29
4 The 1988 Basal Stem Rot Incidence (%) on North Estate Segregated According to Former Crops and Age Groups .....	32
5 The 1988 Basal Stem Rot Incidence (%) on South Estate Segregated According to Former Crops and Age Groups .....	33
6 The 1988 Basal Stem Rot Incidence (%) on East Estate Segregated According to Former Crops and Age Groups .....	34
7 The 1988 Basal Stem Rot Incidence (%) on West Estate Segregated According to Former Crops and Age Groups .....	35
8 Influence of Replanting Techniques on the Incidence of Basal Stem Rot in Oil Palm at 15 Years from Field Planting ...	37
9 Relationship of Soil Types with Incidence of Basal Stem Rot on Palms Greater than 11 Years .....	40
10 Influence of Age of Oil Palms on Incidence of Basal Stem Rot in 12 Estates .....	42
11 Source and Origin of <u>Ganoderma</u> Isolates ....	48



Table		Page
12	Characteristics of Substrates Used for <u>Ganoderma</u> Sporophore Induction .....	51
13	Effect of Substrates on the Eg-01 Isolate after Three Weeks of Incubation ....	60
14	Sporophore Development on Four Substrates which Achieved Full Mycelial Colonisation after Five Weeks of Incubation .....	62
15	Substrate Colonisation by Eight <u>Ganoderma</u> Isolates after Three Weeks of Incubation .....	69
16	Basidiospore Morphology of <u>Ganoderma</u> Isolates Raised <u>In Vitro</u> .....	85
17	Effect of Inoculum Substrates on Vegetative Parameters of Oil Palm Seedlings .....	103
18	Effect of Inoculum Substrates on Vegetative Parameters of Oil Palm Seedlings .....	104
19	Effect of Inoculum Substrates on Vegetative Parameters of Oil Palm Seedlings .....	105
20	Systemic Compounds Evaluated in the <u>In Vitro</u> Fungicide Screening Trial .....	118
21	Systemic Compounds Evaluated in the <u>In Vivo</u> Fungicide Screening Trial .....	121
22	Ranking of Foliar Status in the <u>In Vivo</u> Fungicide Screening Trial .....	123
23	Comparative Efficacy of Systemic Fungicides Tested <u>In Vivo</u> Against <u>Ganoderma boninense</u> .....	125

Table		Page
24	Mortality Rate (%) of Treated Palms in the <u>In Vivo</u> Fungicide Screening Trial at Various Assessment Intervals .....	127
25	Results of Assessment on Palm Survivability at 27 Months from Trial Commencement .....	129
26	Results on the Magnesium Analyses of Frond 17 in Surviving Palms at 27 Months from Trial Commencement .....	130
27	Manuring Schedule of Oil Palm Seedlings in the First Pathogenicity Trial .....	148
28	Composition of Cooper's Nutrient Solution in the Second Pathogenicity Trial .....	149
29	Analysis of Variance for Bunch Counts Conducted at 27 months from Trial Commencement in the <u>In Vivo</u> Fungicide Screening Trial .....	150
30	Analysis of Variance for Foliar Status Assessed at 27 Months from Trial Commencement in the <u>In Vivo</u> Fungicide Screening Trial .....	150

## LIST OF FIGURES

Figure		Page
1	Map of Peninsular Malaysia Showing Locations of the 60 Estates Used in the Assessment .....	21
2	Map of Peninsular Malaysia Showing Locations of the 16 Estates with Significant Incidence of BSR .....:.....	24
3	Relationship between Equivalent Hectarage Lost and Expenditure for Census and Control for 1987 and 1988 .....	30
4	Schematic Diagram of Dimension Measurements of the <u>Ganoderma</u> Sporophore. (a-b) = Length; (c-d) = Width; (e-f) = Thickness; (g-h) = Length of Stipe .....	56
5	Mycelial Growth Rates of Eight <u>Ganoderma</u> Isolates .....	68
6	Schematic Diagram of Establishment of Oil Palm Seedlings on the Fungal-seeded or Non-seeded Substrates in the First Trial A = Rubber Wood; B = Mesocarp Fibre of Oil Palm .....	96
7	Schematic Diagram of Establishment of Oil Palm Seedlings on the Fungal-seeded or Non-seeded Substrates in the Second Trial .....	100



## LIST OF PLATES

Plate		Page
1	Spawn of <u>Ganoderma</u> Isolates (Eg-01, Eg-03, Cg-02 and Cg-05) Established on Whole Padi Grains .....	49
2	Heat-resistant Polypropylene Bags Filled with Substrate Complete with Polyvinyl Chloride (PVC) Necks and Cotton Plugs after Five Weeks of Inoculation .....	53
3	Growth Chamber for Induction of Sporophores of <u>Ganoderma</u> Isolates .....	54
4	Effect of Basal Substrates on Mycelial Growth of Eg-01 <u>Ganoderma</u> Isolate. MFOP (PPF) = Mesocarp Fibre of Oil Palm; SD = Sawdust; WAI = Weeks after Inoculation or Incubation .....	61
5	Early stages of Sporophore Development of Eg-01 Isolate at 35 Days after Incubation in the Growth Chamber .....	64
6	Sporophore Development of Eg-01 Isolate at 44 Days after Incubation in the Growth Chamber .....	65
7	Sporophore Development at 52 Days after Incubation. Note the Brown Colour Deposits of Basidiospores on the Surface of Sporophores .....	67
8	Sporophores of Eg-01, from Dusun Durian Estate, Banting, Selangor, Growing on Mesocarp Fibre of Oil Palm with Addition of Limestone Dust at 5% w/w and Soybean Juice at 32% v/w .....	71



Plate		Page
9	Sporophores of Eg-03, from Dusun Durian Estate, Banting, Selangor, Growing on Mesocarp Fibre of Oil Palm with Addition of Limestone Dust at 5% w/w and Soybean Juice at 32% v/w .....	72
10	Sporophores of Eg-104, from Lian Hup Estate, Muar, Johore, Growing on Mesocarp Fibre of Oil Palm with Addition of Limestone Dust at 5% w/w and Soybean Juice at 32% v/w .....	74
11	Sporophores of Eg-107, from Lian Hup Estate, Muar, Johore, Growing on Mesocarp Fibre of Oil Palm with Addition of Limestone Dust at 5% w/w and Soybean Juice at 32% v/w .....	75
12	Sporophores of Cg-02 (right) Obtained from Dead Coconut, Teluk Intan, Perak in Comparison with an Oil Palm Isolate Eg-01 (left) .....	77
13	Sporophores of Gl-115 or <u>G. lucidum</u> , a Medicinal Fungus, Growing on Mesocarp Fibre of Oil Palm with Addition of Limestone Dust at 5% w/w and Soybean Juice at 32% v/w .....	78
14	Scanning Electron Micrographs of Basidiospores of Eg-01, an Oil Palm Isolate Obtained from Dusun Durian Estate in Selangor .....	80
15	Scanning Electron Micrographs of Basidiospores of Eg-104, an Oil Palm Isolate Obtained from Lian Hup Estate in Johore .....	81
16	Scanning Electron Micrographs of Basidiospores of Gl-115 or the Medicinal <u>G. lucidum</u> .....	82

Plate		Page
17	Scanning Electron Micrographs of Basidiospores of Hg-117 or the Rubber Isolate, <u>G. philippii</u> .....	83
18	Three Distinct Colour Groupings of <u>Ganoderma</u> Isolates, viz., Eg-01 (top), Eg-104 (bottom right) and Gl-115 (bottom left) .....	90
19	Sporophores of the Rubber Isolate, Hg-117 or <u>G. philippii</u> .....	92
20	Rubber Wood Serving as Substrate for <u>G. boninense</u> Inoculum. Inoculated Substrate was Prepared and Incubated in the Dark at 25 + 2°C for Ten Weeks before Use .....	95
21	Polypropylene Bags Enclosed in Inverted Black Polythene Bags Placed in Open Boxes with Compartments and Raised under 50% Overhead Shade .....	101
22	Progressive Dessication of Leaves from Oldest to Youngest is Typical Symptom in Infected Seedlings .....	107
23	<u>Ganoderma</u> Sporophore Formed at the Base of Seedlings where the Pathogen was Inoculated on Rubber Wood as Substrate. Four out of Ten Seedlings Produced this Symptom .....	108
24	Different Stages of Infection by <u>G. boninense</u> on Seedlings in the Types of Substrate Trial at 11 Months. Note that Death has Occurred in Two Seedlings .....	109

Plate		Page
25	Severe Root Rotting (right) of Seedlings which had Displayed Foliar Symptoms. Healthy Roots (left) on Seedlings Inoculated with Inert Rubber Wood .....	110
26	Decay of Stem when Infection has Reached the Stage of a Few Lower Frond Dessication (right) as Compared to a Healthy Seedling (left) .....	111
27	Comparison of a Stunted Seedling on Mesocarp Fibre of Oil Palm Substrate Inoculated with <u>G. boninense</u> (right) with a Normal Growing in the Check Uninoculated Counterpart (left) .....	113
28	Comparison of a Seedling Infected by <u>G. boninense</u> (left) with a Healthy Specimen (right) .....	114
29	Application of Fungicide by Soil Drenching Using a Watering Can. Note that Frond Butts Removal of Trial Palms is a Pre-treatment Selection Procedure .....	122



Abstract of thesis submitted to the Senate of Universiti  
Pertanian Malaysia in fulfilment of the requirements for the degree of  
Master of Agricultural Science.

**BASAL STEM ROT OF OIL PALM :  
INCIDENCE, ETIOLOGY AND CONTROL**

By

**KHAIRUDIN BIN HASHIM**

**July 1990**

Supervisor : Associate Professor Dr. Lim Tong Kwee

Faculty : Agriculture

Basal stem rot (BSR) caused by Ganoderma spp. is a disease of economic importance in oil palm in Malaysia. Although much is known on the occurrence of this devastating disease, fundamental studies on it and its pathogen are rather limited. At present, there appears to be ambiguity in the identity of the causal organisms; there is difficulty in getting oil palms to be artificially infected and that control measures advocated are largely inadequate to contain the disease. Present studies are thus focussed on answering these issues.

An assessment of the BSR situation on selected estates was also carried out to assist in the understanding of the disease.

The assessment carried out showed that serious incidence of the disease was confined to coastal and riverine estates. However, the disease can now be seen to occur in a few inland estates albeit sporadically and at very low incidence. It was demonstrated that previous crops did not necessarily dictate significant incidence of BSR. The underplanting technique and certain soil types can predispose plantings to much higher BSR incidence.

To-date, the fungus has not been cultivated successfully in the laboratory until sporophore formation. A method of cultivation has been developed where this can be achieved under controlled growing conditions. Observation of sporophores has enabled better identification of the pathogen. In that sporophores exhibited polymorphism, the colour of sporophores proved to be a reliable parameter. It has been confirmed that isolates pathogenic to oil palm were not G. lucidum. Two species have been identified in association with diseased palms. They are G. boninense and G. tornatum. An isolate from a dead coconut trunk revealed it as G. boninense while the rubber isolate is G. philippii.



When basidiospores on sporophores were viewed under the scanning electron microscope, it was evident that morphological characters were useful parameters for identification. Apart from the shape and size of basidiospores, the surface ornamentation was a useful tool for taxonomic classification. Differences among G. lucidum, G. philippii, G. boninense and G. tornatum were reinforced.

Previously, doubts have been raised on the pathogenicity of Ganoderma spp. on oil palm. The present study confirmed G. boninense to be a primary parasite. A method of obtaining 100% infection from artificial inoculation has been devised. The Koch's Postulate of duplicating symptoms on very young seedlings similar to those observed on naturally infected field palms and subsequent reisolation of the pathogen were successfully obtained.

In the study on the effects of new systemic fungicides for controlling BSR, it was found that fungicides evaluated in vitro were highly inhibitory to the pathogen. The results showed that hexaconazole, cyproconazole and triadimenol were the most inhibitory to G. boninense with ED<sub>50</sub> values of 0.030, 0.043 and 0.060 ppm a.i. respectively. Propiconazole, fusilazol, carboxin, tridemorph, triadimefon and flutriafol were the next effective with ED<sub>50</sub> values



within the range of 0.2 to 0.6 ppm a.i. The least inhibitory was oxycarboxin with an ED<sub>50</sub> value of 11.9 ppm a.i. As for in vivo fungicide screening, seven fungicides tested were effective at delaying death of infected palms significantly as compared to untreated checks and those treated by surgery. They were fusilazol, hexaconazole, cyproconazole, flutriafol, triadimenol, tridemorph and oxycarboxin in descending order of efficacy.



Abstrak tesis yang dikemukakan kepada Senat Universiti Pertanian Malaysia sebagai memenuhi syarat untuk Ijazah Master Sains Pertanian.

**PENYAKIT REPUT PANGKAL BATANG PADA KELAPA SAWIT:  
INSIDEN, ETIOLOGI DAN KAWALANNYA**

Oleh

**KHAIRUDIN BIN HASHIM**

**Julai 1990**

Penyelia : Professor Madya Dr. Lim Tong Kwee  
Fakulti : Pertanian

Penyakit reput pangkal batang yang disebabkan oleh Ganoderma spp. adalah sejenis penyakit yang penting dari segi ekonomi pada tanaman kelapa sawit di Malaysia. Sungguhpun banyak yang telah diketahui mengenai penyakit yang membinasakan ini, kajian asas tentangnya dan patogen penyebabnya adalah terhad. Pada masa kini, masih terdapat keraguan mengenai identiti organisma penyebab; kesukaran untuk mendapatkan jangkitan buatan pada kelapa sawit dan langkah-langkah pengawalan yang disyorkan kebanyakan kurang memuaskan bagi mengawal merebaknya penyakit ini. Fokus kajian-kajian kini adalah untuk mencari

jawapan bagi isu-isu yang disebutkan. Satu penilaian mengenai situasi penyakit reput pangkal batang di ladang-ladang juga dijalankan bagi membantu dalam memahami penyakit ini.

Penilaian yang dijalankan telah menunjukkan bahawa insiden penyakit adalah terbatas pada ladang-ladang yang terletak di tepi-tepi pantai dan sungai. Bagaimanapun, penyakit ini telah mula dilihat berlaku di beberapa buah ladang yang terletak di kawasan jauh dari laut tetapi insidennya adalah jarang dan rendah. Adalah didapati bahawa tanaman terdahulu tidak semestinya menyebabkan insiden penyakit yang bermakna. Kaedah penanaman semula kelapa sawit baru di bawah pohon tua dan sesetengah jenis tanah boleh mendedahkan tanaman tersebut kepada insiden penyakit reput pangkal batang yang lebih tinggi.

Sehingga kini, kulat penyebab penyakit ini belum berjaya dipelihara di dalam makmal hingga ke peringkat pembentukan sporofor. Satu kaedah pemeliharaan telah direka di mana ini boleh dicapai di bawah keadaan pertumbuhan yang terkawal. Pengamatan ke atas sporofor-sporofor telah membolehkan pengenalan patogen dengan baik. Sporofor didapati mempamerkan pelbagai bentuk atau polimorfisme manakala warnanya merupakan ciri yang berguna untuk pengenalan. Pencilan-pencilan yang patogenik terhadap kelapa sawit telah dikenalpasti bukannya G. lucidum. Dua spesies telah dikenalpasti

berhubung-kait dengan palma berpenyakit. Mereka adalah G. boninense dan G. tornatum. Satu pencilan yang diambil dari batang kelapa yang sudah mati menunjukkan ia adalah G. boninense manakala pencilan dari getah ialah G. philippii.

Apabila basidiospora atas sporofor dilihat di bawah mikroskop elektron pengimbasan, adalah didapati sifat-sifat morfologi berguna untuk pengenalan. Selain dari bentuk dan saiz basidiospora, perhiasan permukaan adalah berguna untuk pengelasan taksonomi. Perbezaan di antara G. lucidum, G. philippii, G. boninense dan G. tornatum diperkukuhkan.

Sebelum ini, keraguan telah dibangkitkan mengenai kepatogenan Ganoderma spp. terhadap kelapa sawit. Kajian ini telah mengesahkan bahawa G. boninense adalah parasit utama. Satu kaedah untuk mendapatkan jangkitan 100% dari inokulasi buatan telah dimajukan. Dalil Koch untuk mendapatkan simptom ke atas anak benih muda kelapa sawit seperti yang terdapat pada pohon kelapa sawit di ladang secara asli serta pengasingan semula patogen telah dilakukan dengan jayanya.

Di dalam kajian ke atas kesan racun-racun kulat sistemik yang baru bagi pengawalan penyakit, adalah didapati bahawa kesemua racun kulat adalah sungguh berkesan in vitro terhadap patogen. Keputusan

menunjukkan bahawa 'hexaconazole', 'cyproconazole', dan 'triadimenol' adalah yang paling berkesan terhadap G. boninense dengan nilai  $ED_{50}$  0.030, 0.043 dan 0.060 bsj b.a. 'Propiconazole', 'fusilazol', 'carboxin', 'tridemorph', 'triadimefon' dan 'flutriafol' adalah juga berkesan dengan nilai  $ED_{50}$  di antara 0.2 hingga 0.6 bsj b.a. Racun kulat yang paling kurang berkesan ialah oxycarboxin dengan nilai  $ED_{50}$  11.9 bsj b.a. Berhubung dengan saringan in vivo racun kulat, tujuh darinya didapati berkesan melengahkan kematian pohon kelapa sawit dengan bermakna berbanding dengan kawalan dan pohon-pohon sawit yang diperlakukan surgeri. Racun-racun kulat tersebut ialah 'fusilazol', 'hexaconazole', 'cyproconazole', 'flutriafol', 'triadimenol', 'tridemorph' and 'oxycarboxin' dalam susunan keberkesanan yang berkurangan.



## CHAPTER I

### INTRODUCTION

The oil palm (Elaeis guineensis Jacq.) is a native of West Africa and was introduced into Malaysia around 1875. The first commercial planting began in 1917 in Selangor, but it was not until the 1960s that the area began to expand rapidly (Hartley, 1977). Within the last three decades, area under oil palm has increased to 1,685,600 hectares in 1987 from about 55,000 hectares in 1960. Production of crude palm oil has also increased from around 82,000 tonnes to 4,532,000 tonnes over the same period (Malaysian Ministry of Finance, 1988). Presently, Malaysia is the leading producer of palm oil in the world, with an estimated total palm product export earnings of M\$3.9 billion in 1987 (Palm Oil Registration and Licensing Authority, 1988).

Although relatively young, the oil palm industry has reflected the success of Malaysia's agricultural diversification programme to avoid over-dependance on rubber as the sole export crop. Major contributing factors for this success are: the politically stable climate for development, the extremely suitable weather with fairly uniform rainfall, the particularly suitable soils with good physical and chemical characteristics, the development of high yielding planting materials through research and development efforts, and lastly, freedom

